

Trientine tetrahydrochloride (Cuprior)

Presentation by Rupert Purchase
(Visiting Fellow, University of Sussex)

BASL Wilson's Disease Special Interest Group Meeting

Friday 29th November 2019

Queen Square House, Institute of Neurology, UCL



Cuprior 150 mg film-coated tablets



gmporphan
*drugs
for orphan
diseases*


Triethylenetetramine tetrahydrochloride (trientine tetrahydrochloride; Cuprior™)

- First reported in 1920
- Crystalline solid mp 266-270 °C
- Prepared by addition of HCl to triethylenetetramine
- Not reported to be hygroscopic
- Cu^{2+} coordination noted in 1936




Trientine (triethylenetetramine) structures


trientine free base (Tosoh, Japan)*
cost (technical grade): ~ 0.0006p/mg



Cufence (Univar: trientine.2HCl)
200 mg free base/capsule
cost/capsule: 15p/mg (free base)



Cuprior (GMP-Orphan: trientine.4HCl)
150 mg free base/tablet
cost/tablet: 25p/mg (free base)



*Quotation from Henan Tianfu Chemical Co., China

Cuprior 150 mg tablets

- Stable at room temperature; do not require refrigeration
- 16 mm x 8 mm – each tablet dividable into equal doses
- Recommended adult dose: 3–6½ tablets/day
- 6 x Cuprior tablets/day \equiv 0.9 grams chloride ion

[Dietary chloride ion intake = 2 grams/day]

Cuprior: 150 mg base \equiv 300 mg trientine.4HCl; Cufence: 200 mg base \equiv 300 mg trientine.2HCl

GMP-Orphan – EMA submission data (2015)

Cuprior tablets compared with Cufence capsules

Dissolution

- Faster aqueous dissolution (pH 1.2; pH 4.0 and pH 6.8)
- Ascribed to differences in formulation rather than solubility differences between the two salts

Bioavailability

- Higher bioavailability *cf.* Cufence (pharmacokinetic data)
- Approx. 60% dose adjustment factor proposed to provide daily dose of trientine free base equivalent to that from the dihydrochloride

Bioavailability of Cuprior and Cufence

Relative bioavailabilities determined by:

- Aqueous solubility and dissolution
- Protonation of trientine base (bis-protonated at pH 6-7; tetra-protonated at pH 2-3)
- pH of gastric acid (pH 1.5 to 3.5)
- pH of intestinal tract (pH 6 to 7.4)

Bioavailability of Cuprior and Cufence

Conclusions:

- The speciation of trientine *in vivo* will be the same irrespective of the administration of the dihydrochloride or the tetrahydrochloride
- Once in the intestinal tract, the subsequent absorption, distribution and metabolism of the two drugs will also be equivalent

Therefore:

Relative bioavailability of trientine is determined by aqueous solubility/dissolution of the formulated salt

Cuprior: clinical trial

Trientine tetrahydrochloride (TETA.4HCl) for the treatment of Wilson's disease

- ClinicalTrials.gov Identifier: NCT03539952
- Participating countries: Belgium, Brazil, Denmark, France, Germany, Italy, Poland, Sweden, UK and USA
- Completion date: November 2021

<https://www.clinicaltrials.gov/ct2/show/NCT03539952>